

Taro Yamane: Sample Size Table

Table 1 **Sample Size for $\pm 3\%$, $\pm 5\%$, $\pm 7\%$, $\pm 10\%$, Precision Levels where Confidence level is 95%**

Size of Population	Sample Size (n) for Precision (e) of:			
	$\pm 3\%$	$\pm 5\%$	$\pm 7\%$	$\pm 10\%$
500	a	222	145	83
600	a	240	152	86
700	a	255	158	88
800	a	267	163	89
900	a	277	166	90
1,000	a	286	169	91
2,000	714	333	185	95
3,000	811	353	191	97
4,000	870	364	194	98
5,000	909	370	196	98
6,000	938	375	197	98
7,000	959	378	198	99
8,000	976	381	199	99
9,000	989	383	200	99
10,000	1,000	385	200	99
15,000	1,034	390	201	99
20,000	1,053	392	204	100
25,000	1,064	394	204	100
50,000	1,087	397	204	100
100,000	1,099	398	204	100
>100,000	1,111	400	204	100

a = Assumption of normal population is poor (Yamane, 1967). The entire population should be sampled.

Table 2 Sample Size for $\pm 5\%$, $\pm 7\%$, $\pm 10\%$, Precision Levels where Confidence level is 95%

Size of Population	Sample Size (n) for Precision (e) of:		
	$\pm 5\%$	$\pm 7\%$	$\pm 10\%$
100	81	67	51
125	96	78	56
150	110	86	61
175	122	94	64
200	134	101	67
225	144	107	70
250	154	112	72
275	163	117	74
300	172	121	76
325	180	125	77
350	187	129	78
375	194	132	80
400	201	135	81
425	207	138	82
450	212	140	82